

Exercise 1.1. [Short Questions]

Let M be a set of formulas, and let F and G be formulas. Which of the following assertions hold?

1. If F satisfiable then $M \models F$ $F = A_1 \quad M = \{A_2\}$

2. F is valid iff $\top \models F$

3. If $\models F$ then $M \models F$

4. If $M \models F$ then $M \cup \{G\} \models F$

5. $M \models F$ and $M \models \neg F$ cannot hold simultaneously $M = \{\perp\}$

6. If $M \models G \rightarrow F$ and $M \models G$ then $M \models F$

$M \models F$ iff for all A for all $M_i \in M$, $A \models M_i$; implies $A \models F$

iff for all A for all $M_i \in M$, $A \models M_i$; implies $A \models (G \rightarrow F) \wedge G$

iff for all A for all $M_i \in M$, $A \models M_i$; implies $A \models G \rightarrow F$ and $A \models G$

iff for all A for all $M_i \in M$, $A \models G \rightarrow F$ and for all $M_i \in M$, $A \models G$

iff $M \models G \rightarrow F$ and $M \models G$

Exercise 1.2. [Coincidence Lemma]

Assume that for all atomic formulas A_i in F , $\mathcal{A}(A_i) = \mathcal{A}'(A_i)$. Show that

$$\mathcal{A} \models F \text{ iff } \mathcal{A}' \models F$$

Induction over structure \models

- Case $\models = A$; for some A :
 $A \models A$; iff $\mathcal{A}(A_1) = 1$ iff $\mathcal{A}'(A_1) = 1$ iff $A' \models A$,
- Case $\models = \neg G$ for some G . Assume $A \models G$ iff $\mathcal{A}' \models G$. (IH)
 $A \models \neg G$ iff $A \not\models G$ iff $\mathcal{A}' \not\models G$ (by IH)
 iff $\mathcal{A}' \models \neg G$
- Case $\models = G \vee H$. Assume $A \models G$ iff $\mathcal{A}' \models G$
 and $A \models H$ iff $\mathcal{A}' \models H$
 $A \models G \vee H$ iff $A \models G$ or $\mathcal{A}' \models H$
 iff $\mathcal{A}' \models G$ or $\mathcal{A}' \models H$ (by IH)
 iff $\mathcal{A}' \models G \vee H$
- Case $\models = G \wedge H$ analogous

Exercise 1.3. [Semantic Proof]

Let $\models F \rightarrow G$ where F and G do not share any atoms. Show that then either F is unsatisfiable or G is a tautology (or both). Hint: you may want to use the previous result.

$$\models F \rightarrow G \quad \text{iff} \quad \models \neg F \vee G$$

||

$$\neg F \odot G$$

$$\models F \rightarrow G$$

$$\begin{aligned} & \text{iff} \quad \models \neg F \vee G \\ & \text{(Let } A_F : M_F \rightarrow \{0,1\} \text{ and } A_G : M_G \rightarrow \{0,1\} \text{ atoms appearing in } F) \\ & \quad \text{Let } A_F \vdash \neg F \text{ and } A_G \vdash G \text{ (satisfy, satisfy)} \end{aligned}$$

$$\text{iff for all } A, A \models \neg F \vee G$$

$$\text{iff for all } A, A \models \neg F \text{ or } A \models G$$

$$\text{iff for all } A, A \models \neg F \text{ or } A \models G \text{ or } A \models \neg F \wedge A \models G$$

$$\text{iff for all } A, A \models \neg F \text{ or } A \models G$$

$$\text{iff for all } A, A \models \neg F \text{ or for all } A, A \models G$$

$$\text{iff for all } A, A \models \neg F \text{ or for all } A, A \models G$$

$$\text{iff } \models \neg F \text{ or } \models G$$

Exercise 1.4. [Satisfiability Algorithms]

Check the following formulas for satisfiability using one of the algorithms seen in the lecture:

$$\neg(A \wedge D)$$

- $F_1 = (\neg A \vee \neg D \vee B) \wedge D \wedge \neg B \wedge E \wedge (\neg D \vee \neg E \vee C)$

How to Form. (c) CNF

No clause like $A \vee B$

→ Add mark one positive literal in each clause

$$F_1 \equiv (A \wedge D \rightarrow B) \wedge D \wedge \neg B \wedge E \wedge (\neg D \wedge E \rightarrow C)$$

$$\wedge (T \rightarrow D) \wedge (B \rightarrow T) \wedge (T \rightarrow E)$$

Init Place

Mark D, E

Mark C.

RETURN N SAT

$$A(B) = A(A) = 0$$

$$A(C) = A(D) = A(E) = 1$$

Satisfying Assignment

$$F_2 = (A \rightarrow C) \wedge (C \rightarrow E) \wedge (\top \rightarrow B) \wedge (C \wedge A \rightarrow D) \wedge (B \rightarrow A) \wedge (B \rightarrow E) \wedge (D \wedge E \rightarrow \perp)$$

Mark P init

Mark A, E

Mark C

Mark D

Return UNSAT

$$F_3 = (A \rightarrow E) \wedge (B \rightarrow \perp) \wedge (C \rightarrow B) \wedge (\top \rightarrow A) \wedge (E \wedge B \rightarrow C) \wedge (C \rightarrow D)$$

Mark A $\boxed{A(A) = A(E) = 1}$ ← Any model has to contain this
Mark E $\boxed{A(B) = A(C) = A(D) = 0}$

Return SAT

↑
For the unmarked variables there may be variability,
but the algorithm ensures that this assignment is satisfying
if it returns "SAT".